

Unified approach to redshift in cosmological /black hole spacetimes and synchronous frame

A. V. Toporensky

*Sternberg Astronomical Institute, Lomonosov Moscow State University,
Universitetsky Prospect, 13, Moscow 119991, Russia and
Kazan Federal University, Kremlevskaya 18, Kazan 420008, Russia**

O. B. Zaslavskii

*Department of Physics and Technology,
Kharkov V.N. Karazin National University,
4 Svoboda Square, Kharkov 61022, Ukraine and
Kazan Federal University, Kremlevskaya 18, Kazan 420008 Russia†*

S. B. Popov

*Sternberg Astronomical Institute, Lomonosov Moscow State University,
Universitetsky Prospect, 13, Moscow 119991, Russia*

Usually, interpretation of redshift in static spacetimes (for example, near black holes) is opposed to that in cosmology. In this methodological note we show that both explanations are unified in a natural picture. This is achieved if considering the static spacetime one (i) makes a transition to a synchronous frame, (ii) returns to the original frame by means of local Lorentz boost. To reach our goal, we consider a rather general class of spherically symmetric spacetimes. In doing so, we construct frames that generalize the well-known Lemaitre and Painlevé–Gullstrand ones and elucidate the relation between them.

PACS numbers: 04.20.-q; 04.20.Cv; 04.70.Bw

*Electronic address: atopor@rambler.ru

†Electronic address: zaslav@ukr.net

I. INTRODUCTION

Though calculation of redshifts in General Relativity (GR) has no principal difficulties, their interpretation continues to be a source of debates. It is well known that gravitational field can induce a blue/red shift which is absent in Special Relativity (SR). For example, a photon emitted near a black hole horizon reaches a distant observer severely redshifted even in the case where emitter and observer do not move with respect to each other. As far as the gravitational redshift of this type is concerned, it can be obtained as follows. In the static field one can introduce such a time variable t that different static observers measures the same time intervals dt between a pair of corresponding events (say, emission and absorption of photons). Then, the intervals of the proper time differ by the ratio of the factors $\sqrt{g_{tt}}$, where g_{tt} is the corresponding component of the metric tensor. One can also describe this redshift in a more formal fashion. In the static gravitation field there exists a timelike Killing vector ξ^μ . Let a photon have the wave vector k^μ . Then, the quantity $\omega_0 = -k_\mu \xi^\mu$ is conserved along the path of a photon due to the geodesic equation for k^μ and Killing equations. It has the meaning of a frequency measured at infinity. Let us imagine a set of static observers with the four-velocities u^μ . They measure the local frequencies $\omega = -k_\mu u^\mu$. For a static observer $u^\mu = \xi^\mu/N$, where $N = \sqrt{-\xi_\mu \xi^\mu}$ is a lapse function. Then, we obtain that $\omega N = \omega_0$ which gives a redshift in a static field. These two methods imply staticity of the field and are completely equivalent.

However, the cosmological redshift does not fall into this scheme. In an expanding universe, one is tempted to recourse to the explanation in the spirit of the Doppler effect. Meanwhile, it cannot be simply reduced to this effect (at least globally, see discussion below), since, in particular, coordinate recession velocities can exceed the speed of light, which is absolutely impossible in SR; so gravitational field plays an essential role. However, we cannot calculate the redshift as it is done in the black hole spacetime because in expanding or contracting Universe there is no timelike Killing vector. Gravitational time delay is also absent in the Friedmann-Robertson-Walker (FRW) coordinate system because this system is synchronous, which means that all particles in the Hubble flow share the same time that coincides with the proper time. Due to this reason, the cosmological redshift is sometimes treated as the third type of redshift which is distinct from both the Doppler and gravitational ones (see, e.g. Sec. 15 of [1]).

Such classification of redshifts, being rather useful from a practical point of view (since it prevents from using incorrect formulae for inappropriate types of redshifts), has its own conceptual problems. The reason is that in this picture cosmology appears to be separated from other GR issues which seems rather strange, and what is more important, such division cannot be done technically in many situations, which are slightly more sophisticated than the FRW metrics. For example, do redshifts in the Schwarzschild-de Sitter solution might be interpreted in cosmological terms, or it is better to apply terminology from black hole physics?

The goal of the present methodological paper is to show that apparent “cosmological” features of cosmological redshift are not limited only by situations related to cosmology, but, rather, follow from properties of synchronous coordinate system (for example, the FRW metric), and, so can be present in any other applications of GR (including black holes) when an appropriate coordinate system is chosen.

There are also other unifying approaches to redshifts. One of them is based on parallel transport of the observer’s velocity along the lightlike geodesic [2], [3], [4]. By contrast, we do not introduce additional constructions and operate with equations of motion directly.

II. COSMOLOGICAL REDSHIFT

Derivation of the cosmological redshift is the case of controversy — different authors follow different routes to obtain this quantity. We briefly mention the most popular of them, and then present another one, which will be used below.

We start with the usual expression for the interval (only radial motions are discussed):

$$ds^2 = -c^2 dt^2 + a^2(t) d\chi^2. \quad (1)$$

Here c is the light speed.

The simplest way to obtain the redshift in the FRW metric is to use the conformal time, η . Defining $\eta(t) = \int dt/a$ we reduce the metric to the form

$$ds^2 = a^2(d\chi^2 - d\eta^2). \quad (2)$$

Assuming $\chi_{obs} = 0$ we obtain $\chi_{em} = \eta_{obs} - \eta_{em}$. Here, subscripts “obs” and “em” refer to an observer and emitter, respectively. If a signal was emitted during a period $\Delta\eta$, then it

is observed during the same interval of conformal time. However, physical time intervals at two moments are different: $\Delta t_{em} = a(t_{em})\Delta\eta$ and $\Delta t_{obs} = a(t_{obs})\Delta\eta$. Thus, $\Delta t_{em}/\Delta t_{obs} = a(t_{em})/a(t_{obs})$, where we neglected the change of the scale factor during the processes of emission and observation.

Then, one can use symmetry of a metric to derive the redshift using conservation laws (see pp. 457-45 of [5]). It is possible to show that in the FRW metric the momentum changes as $a(t)^{-1}$. Then, as for photons momentum is $\sim \lambda^{-1}$, we obtain that $\lambda \sim a(t)$. Thus, we derive the usual expression for the redshift.

Both of these methods exploit properties of a metric (in particular, FRW), and so they are not general derivations.

The third way we want to mention is popular, however, it is not fundamental, as *a priori* it is not obvious that it might provide the correct answer. This method uses integration of the local Doppler effect along the trajectory of a photon, or even just the statement, that locally cosmological redshift can be given by the Doppler formula. Such a simplified approach, most often with appropriate comments that this is not an accurate – but just illustrative, – way of derivation, can be found in some textbooks (for example, Ch. 5.2 of [6]).

The most general method one can often find in textbooks uses null geodesics (see pp. 777 - 778 of [5]).

For photons

$$ds^2 = 0. \tag{3}$$

Thus, we obtain from (1) the expression for the coordinate velocity:

$$d\chi/dt = c/a.$$

Then, let us consider two pulses of an electromagnetic wave. They are emitted in two moments of time t_{em1} , t_{em2} and observed, correspondingly, at t_{obs1} and t_{obs2} . Both are emitted by a galaxy at χ_{em} . The observer is situated at χ_{obs} .

If an emitter and an observer follow the geodesic lines, their values χ remain fixed since (1) represents a synchronous system. World lines of the two photons are:

$$\chi_{em} - \chi_{obs} = c \int_{t_{em1}}^{t_{obs1}} a^{-1} dt, \quad (4)$$

$$\chi_{em} - \chi_{obs} = c \int_{t_{em2}}^{t_{obs2}} a^{-1} dt. \quad (5)$$

Then

$$\int_{t_{em1}}^{t_{obs1}} a^{-1} dt - \int_{t_{em2}}^{t_{obs2}} a^{-1} dt = 0. \quad (6)$$

Eq. (6) can be rewritten in the equivalent form

$$\int_{t_{obs1}}^{t_{obs2}} a^{-1} dt - \int_{t_{em1}}^{t_{em2}} a^{-1} dt = 0. \quad (7)$$

Neglecting changes in a during the interval of emission of the two pulses, and during the interval of their observation (i.e. putting $a(t_{em1}) = a(t_{em2})$ and $a(t_{obs1}) = a(t_{obs2})$), we obtain

$$\frac{t_{obs2} - t_{obs1}}{a(t_{obs})} - \frac{t_{em2} - t_{em1}}{a(t_{em})} = 0.$$

At emission $\lambda_{em} = c(t_{em2} - t_{em1})$. At observation $\lambda_{obs} = c(t_{obs2} - t_{obs1})$. So,

$$\lambda_{em}/a(t_{em}) = \lambda_{obs}/a(t_{obs}).$$

Finally, we come to the usual expression for the cosmological redshift.

In our paper we use a variation of this method. In our opinion, it is, in the first place, interesting from the methodological point of view and also provides a clear visualization of the process. For our goals here, this version of derivation is needed because it, unlike the method above, can be easily generalized to inhomogeneous metrics, which will be used below. It essentially relies on using different definitions of velocity. This contains some subtleties discussed below.

III. DEFINITIONS OF VELOCITY AND “LIGHT FASTER THAN LIGHT”

We note that the fact that the coordinate velocity of light, $d\chi/dt$, can be not equal to c in some coordinate systems is obvious, because coordinates have no direct physical meaning. What is less obvious is that the velocity of light defined as derivative of the proper distance

from an observer to a photon over proper time (which in the FRW metric is equal to the coordinate time) can be different from c as well.

To work with proper distance we need to integrate $\sqrt{g_{\chi\chi}}$ over $d\chi$ at the same moment of proper time. This means that we have to define hypersurfaces $t = \text{const.}$ To do this, it is necessary to fix a coordinate system. Then, the answer depends on the coordinate system, i.e. is not invariant.

In the FRW metric (1) the proper distance between $\chi = 0$ and some χ at a given moment of time t is equal to $r = \chi a$. If χ is allowed to vary with time, for the velocity dr/dt we can write a chain of equations:

$$\frac{dr}{dt} = \frac{d(\chi a)}{dt} = \frac{d\chi}{dt}a + \frac{da}{dt}\chi = c + \frac{da}{dt}\chi = c + \frac{\dot{a}}{a}a\chi = c + v, \quad (8)$$

where $v = Hr$ is the Hubble flow velocity, and $H = \dot{a}/a$ is the Hubble constant. This clearly demonstrates so-called “superluminal motion” in cosmology. Note, that eq. (8) can be considered as a particular case of the general fact that in cosmology peculiar velocity is summed with the Hubble velocity following the Galilean law. It is worth stressing that a local velocity of light defined as $ad\chi/dt$ (the first term in eq. 8) is equal to c according to (3), as it should be.

To make the difference between two definitions of velocity more pronounced, let us denote $P(\chi_2, t \mid \chi_1, t)$ the proper distance between points 1 and 2 at the moment t . Then,

$$\frac{dr}{dt} = \frac{dP(\chi, t \mid 0, t)}{dt} = v_{loc} + v_{fl}, \quad (9)$$

where the local velocity

$$v_{loc} = \lim_{\Delta t \rightarrow 0} \frac{P(\chi + \Delta\chi, t \mid \chi, t)}{\Delta t} = \frac{\partial P(\chi, t \mid 0, t)}{\partial \chi} \frac{d\chi}{dt}, \quad (10)$$

the flow velocity

$$v_{fl} = \frac{\partial P(\chi, t \mid 0, t)}{\partial t}. \quad (11)$$

It is the quantity v_{loc} that can be found by pure local measurements. For light, $v_{loc} = c$. The quantity dr/dt is more involved and cannot be detected by direct local measurements. Not surprisingly, it can exceed c without any contradiction with GR and SR.

Now let us derive the equation for the cosmological redshift. Here we will use the fact that two light pulses, being at slightly different distances from an observer, have different

velocities with respect to this observer. So, let us consider two neighboring maxima of a light wave directed toward an observer, separated by a spatial distance $\Delta r = \lambda$. This radial difference results in the velocity difference with respect to the observer $\Delta v = H\Delta r$. This means that the velocity of the maximum which is closer to the observer is less than the velocity of the maximum which is behind it. Due to this difference, the distance between these maxima changes and we can construct a differential equation

$$d\Delta r/dt = H\Delta r. \quad (12)$$

After obvious calculations using the definition of the Hubble parameter $H = \dot{a}/a$ we obtain that the ratio of wavelengths at t_{em} and the moment of observation, t_{obs} , is equal to the inverse ratio of scale factors at these moments: $\lambda_{em}/\lambda_{obs} = a_{em}/a_{obs}$. This is the standard result for the cosmological redshift.

Below, we want to show how features typical for cosmology can be considered in a rather generic black hole background, if a synchronous system is used. In doing so, we generalize the well-known Lemaitre metric connected to an observer free falling into a black hole. Then, it is clear that the concept of an expanding or contracting space (discussed above in the cosmological context) by itself is not necessary to understand the physics of cosmological evolution.

First of all, the definition of the Hubble flow and peculiar velocity has a straightforward generalization. Indeed, in a synchronous system coordinate lines $x^i = const$ are geodesics (see, e.g. Sec. 11.97 of [8]). Observers with constant spatial coordinates share the same time coordinate which coincides with their proper time. One can define a proper distance between two points at some particular time t and also velocity of one point with respect to another by differentiating the proper distance with respect to t . For simplicity, we will consider spherically symmetric metrics.

Let χ be the radial coordinate of the observed particle and χ_2 changes, while the coordinate χ_1 of the observer remains fixed. Then, we get, using the well-known formula for the integral with varying limits:

$$v = \frac{d}{dt} \int_{\chi_1}^{\chi_2} \sqrt{g_{\chi\chi}} d\chi = v_{fl} + v_{loc}, \quad (13)$$

The velocity appears to be a sum of two components. The first one

$$v_{fl} = \int_{\chi_1}^{\chi_2} \frac{d}{dt} \sqrt{g_{\chi\chi}} d\chi \quad (14)$$

is fully due to non-static character of the metric (and can be considered as an analogue of velocity of the Hubble flow). While the second one

$$v_{loc} = \sqrt{g_{\chi\chi}(\chi_2)} d\chi_2 / dt \quad (15)$$

is non-zero only when the radial coordinate of a particle changes in time, so we can treat it as an analogue of peculiar velocity in cosmology. Eqs. (13) – (15) represent the counterpart of eqs. (8) – (11) in cosmology. Of course, for light locally v_{loc} is always equal to c .

IV. STATIC SPACETIME AND RELATIONSHIP BETWEEN FRAMES: LEMAITRE VS. PAINLEVÉ–GULLSTAND METRICS

In what follows, we consider a static metric in a “cosmological manner”. This includes the black hole spacetime. We show similarity between phenomena in black hole physics and cosmology, and demonstrate that, looking different at the first glance, these processes have the same interpretation in a synchronous metric. We discuss a black hole metric of a rather general form. Without the loss of generality, we take it to be

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{f(r)} + r^2(d\theta^2 + \sin^2\theta d\phi^2). \quad (16)$$

It includes the Schwarzschild metric ($f = 1 - r_g/r$), the Reissner-Nordström one, etc. Extension to the case $g_{00}g_{11} \neq -1$ is straightforward. The event horizon lies at $r = r_g$, so $f(r_g) = 0$. Near the horizon, the metric becomes degenerate. This can be remedied by introducing a new set of coordinates.

Making the coordinate transformations [7]

$$\rho = t + \int \frac{dr}{f\sqrt{1-f}}, \quad (17)$$

$$\tau = t + \int \frac{dr}{f} \sqrt{1-f} \quad (18)$$

one can reduce the metric to the form

$$ds^2 = -d\tau^2 + (1 - f)d\rho^2 + r^2(\rho, \tau)(d\theta^2 + \sin^2\theta d\phi^2). \quad (19)$$

If $f = 1 - r_g/r$, we return to the well-known Lemaitre metric [8]. Therefore, we will call (19) the generalized Lemaitre (GL) frame.

Since the GL metric is synchronous, all free falling radial observers have the constant comoving radial coordinate ρ and share the same proper time. It appears that the proper distance is related with r in a very simple way. Namely, fixing τ one finds from (17) – (19) after simple calculations that the proper distance between points 1 and 2 equals

$$l = \int_{\rho_1}^{\rho_2} d\rho \sqrt{1 - f} = r_2 - r_1. \quad (20)$$

In what follows, we will use r along with the radial coordinate ρ . We can see here some analogue of the cosmological situation with the FRW metric where the proper distance between two points is given by the formula $d = a(\chi_2 - \chi_1)$ which is valid for any observer moving in the Hubble flow.

It is worth noting that for massive particles composing the GL system $E = m$, $\rho = \text{const}$ and thus $v_{fl} = v$, $v_{loc} = 0$. For a photon $|v_{loc}| = 1$.

One can also make one more step and write down the metric in a “mixed” form using a new time variable τ and old radial variable r . Then, eliminating dt we have from (17), (18):

$$ds^2 = -d\tau^2 f + dr^2 + 2drd\tau\sqrt{1 - f} + r^2(d\theta^2 + \sin^2\theta d\phi^2). \quad (21)$$

In the Schwarzschild case, the metric (21) is nothing else but the so-called Painlevé–Gullstrand metric (see, e.g. [9] and references therein). For a generic f , it is natural to call it the generalized Painlevé–Gullstrand metric. It is worth noting that it is metric (21) which was used in [10] to describe “a river model of black hole” as an interesting counterpart of the cosmological concept of “expanding space”. Strange as it may seem, the relationship between the Painlevé–Gullstrand and Lemaitre metrics, to the best of our knowledge, was not noticed before.

In these coordinates, some properties of the metric look rather simple. In particular, the hypersurfaces $\tau = \text{const}$ are flat. Eq. (20) becomes direct consequence of (21). For a massive particle falling along geodesic from the rest at infinity one has $dr/d\tau = -\sqrt{1 - f}$, whence we have for the velocity $v = |dr/d\tau|$:

$$v = \sqrt{1 - f}. \quad (22)$$

If points 1 and 2 move along geodesics, their relative velocity is:

$$v_{12} = v_2 - v_1. \quad (23)$$

For light propagation we have:

$$\frac{dr}{d\tau} = \pm 1 - v, \quad (24)$$

where we put $c = 1$ and v coincides with (22). Here, sign “+” corresponds to propagation outward and “-” corresponds to propagation inward.

V. BEHAVIOR OF WAVELENGTH

Now, by the same method as in the FRW cosmology we derive the redshift of a photon emitted by a free falling particle at r_{em} and absorbed by another free falling particle at r_{obs} . Remember that the redshift in the FRW frame appears because the proper distance between two light pulses changes with time since the speed of light with respect to a distant observer depends on the distance between the pulse and the observer. The same situation appears in the Lemaitre metric, as we will see soon. Our goal is to start with two maxima of light wave separated by $\Delta r = \lambda_1$ and trace the evolution of this separation during light propagation. An observer will see the maxima separated by λ_2 which gives us the redshift via $1 + z = \lambda_2/\lambda_1$. As usual for synchronous systems, the coordinate velocity of light is $v_{fl} + c$. For two points with radial difference Δr at r_{em} (here the dependence on radial coordinate at emission is important, because the metric is not homogeneous) corresponding velocities of light differ by $\Delta v_{fl} = v'_{fl}(r)\Delta r$. In this Section, we consider free-falling particles comoving with respect to the GL system, so peculiar velocities vanish and $v_{fl} = v$.

It is worth noting that while we need to subtract the velocity of the observer in order to get relative velocity (23), this does not affect the *difference* between velocities of nearby points Δv , so we can consider the position r_0 of the observer as arbitrary. For simplicity, we consider free-fall velocities with respect to an infinitely distant observer.

So, if at emission the distance between two wave maxima was $|\Delta r| \equiv \lambda$, it later increases according to

$$d\Delta r = \Delta v d\tau = v'(r)\Delta r d\tau. \quad (25)$$

This equation is an analogue of the cosmological equation (12). It is useful to integrate over the radial coordinate r (instead of the Lemaitre coordinates ρ and τ). Let us consider an inwardly moving photon, so that according to (24), $d\tau = -dr/(1+v)$. Then we get:

$$\frac{d\lambda}{dr}(1+v) = \frac{dv}{dr}\lambda, \quad (26)$$

so we have the differential equation for λ .

$$\frac{d\lambda}{\lambda} = \left(\frac{dv}{dr}dr\right)/(1+v). \quad (27)$$

If dv/dr is not a function of time (as it is in the static spherically symmetric spacetime) we can write:

$$\frac{d\lambda}{\lambda} = \frac{dv}{1+v}. \quad (28)$$

Note, that in the FRW cosmology dv/dr is not a function of time only in the de Sitter universe where $dv/dr = H = \text{const}$.

Since for the photon moving inward the sum $v+1$ never crosses zero, the differential equation is regular at the event horizon, so the solution should give us the redshift of such a photon for both possible locations of an emitter and an observer (inside or outside the event horizon).

Then, we have the ratio of the initial and final wavelengths for an inward motion of a photon:

$$\frac{\lambda_{obs}}{\lambda_{em}} \equiv 1+z = \frac{1+v_{obs}}{1+v_{em}}. \quad (29)$$

For the outward motion of a photon, in a similar way one would obtain:

$$1+z = \frac{1-v_{obs}}{1-v_{em}}. \quad (30)$$

These results are in agreement with those of [11] and [12] (in the latter case they were derived with the help of the Kruskal coordinates). As noted above, in FRW cosmology these formulas are valid only for the de Sitter solution. For other regimes the dependence between the redshift and the velocity at emission (in FRW metrics it is reasonable to set the velocity of observer to zero) is more complicated, see, for example, [13].

Going back to static spacetimes we can use (22) to get the general expression for the redshift through radial coordinate of emission and observation:

$$1 + z = \frac{1 \pm \sqrt{1 - f(r_{obs})}}{1 \pm \sqrt{1 - f(r_{em})}} \quad (31)$$

where the upper sign is used for ingoing photon, and the lower sign for the outgoing photon.

The latter case can be applied if observer and emitter are either both outside the horizon (in this situation evidently $r_{em}(\tau_{em}) < r_{obs}(\tau_{obs})$), or both inside the horizon (and $r_{em}(\tau_{em}) > r_{obs}(\tau_{obs})$). On the contrary, there are no restriction for the mutual locations of emitter and observer for the ingoing photon case.

If, for example, an observer crosses the horizon, $f_{obs} = 0$, then according to (22), $v_{obs} = 1$. So, he/she receives a signal from a remote source at infinity ($v_{em} = 0$) with the redshift $z = 1$.

If the observer approaches the singularity where $r \rightarrow 0$ and $f \rightarrow -\infty$ (like in the Schwarzschild metric where $f = 1 - r_g/r$), $v_{em} \rightarrow \infty$, so the redshift diverges.

It is clear from the very method of derivation that these results are quite general and are valid far beyond GR. It is necessary, of course, that free massive particles follow geodesics. This holds in metric theories of gravity. Also, it is necessary that photons move along null geodesics. This is achieved if the coupling between electromagnetism and gravity is minimal. This second condition is fulfilled in most popular generalizations of GR, though may be violated for some class of theories studied by Horndeski [14].

A. Special case: propagation along the horizon

One particular case of an outward propagation of a photon is still beyond our analysis. Namely, if a photon is emitted exactly at the horizon and further propagates along it, we cannot integrate over r because it is constant for such photons (but the Lemaitre coordinate ρ obviously changes).

Instead, we use directly (25) with Δr replaced with λ and $r = r_g$, whence

$$d\lambda = \lambda v'(r_g) d\tau, \quad (32)$$

and $1 + z = \exp[v'(r_g)(\tau_{obs} - \tau_{em})]$. Using (22) and the fact that $f(r_g) = 0$, $f'(r_g) = 2\kappa$, where κ is the surface gravity, we obtain:

$$1 + z = \exp[\kappa(\tau_{obs} - \tau_{em})]. \quad (33)$$

So, in this case the redshift can be expressed through the difference between emitting and observing time in agreement with the result obtained in [7] by another method. In particular case of Schwarzschild metrics the same problem has been considered earlier using Kruskal coordinates [12].

VI. FREQUENCIES IN DIFFERENT FRAMES

In this section we obtain the same results for redshifts in a more formal way as it is usually done if problems of interpretations are not concerned.

The geodesic equations for radial light propagation in the static frame are the following: (16):

$$\begin{aligned} k^t &= \frac{\omega_\infty}{f}, \\ k^r &= -\omega_\infty, \end{aligned} \quad (34)$$

where for definitiveness we consider the motion in the inward direction only. Here, k^μ is the wave vector, ω_∞ has the meaning of the frequency measured by a remote observer at infinity.

Using (17) and (18), one can find

$$k^\tau = \frac{\omega_\infty}{f}(1 - \sqrt{1-f}), \quad (35)$$

$$k^\rho = -\frac{\omega_\infty}{f\sqrt{1-f}}(1 - \sqrt{1-f}). \quad (36)$$

The frequency measured by a local observer with the four-velocity u^μ is:

$$\omega = -k_\mu u^\mu, \quad (37)$$

where $u^\mu = dx^\mu/d\tau$ is the four-velocity, τ is the proper time.

For the GP observers $\rho = \text{const}$, so $u^\tau = 1$, $u^\rho = 0$. Then, we have from (37) that

$$\omega = k^\tau = \frac{\omega_\infty}{f}(1 - \sqrt{1-f}) = \frac{\omega_\infty}{1 + \sqrt{1-f}}. \quad (38)$$

In terms of the velocity, it can be rewritten as

$$\omega = \omega_\infty \frac{(1-v)}{f} = \frac{\omega_\infty}{1+v}. \quad (39)$$

If light is emitted at ρ_{em} and is absorbed at $\rho_{obs} < \rho_{em}$,

$$1+z \equiv \frac{\omega_{em}}{\omega_{obs}} = \frac{1+v_{obs}}{1+v_{em}} \quad (40)$$

which coincides with (29), as it should be. If $\rho_{obs} > \rho_{em}$, we again obtain (30).

VII. FROM THE GL FRAME TO THE STATIC ONE

In the above analysis we follow the usual cosmological pattern to derive the redshift. We can also use these results to get the standard redshift in the Schwarzschild geometry, which is usually obtained using the notation of gravitational time delay, and is described as a typical example of a gravitational redshift. It concerns an observer and emitter with fixed Schwarzschild radial coordinate r . We want to reproduce it using the above formulae without any references to the gravitational “time delay”.

The above formulas describe redshift (or blueshift) in the GL frame, when frequencies or wavelengths are measured by the corresponding free falling observers who have $\rho = \text{const.}$ Now, we are interested in a redshift in the static frame. To obtain it, it is sufficient to make a boost from the GL frame. It follows from (17), (18) that for an observer or emitter to be fixed at a constant r , it must have $\rho = \tau + \text{const.}$ Then, it moves with the velocity $v = dl/d\tau$ (22) with respect to the GL frame. So, we need to combine the redshift (29) with the Lorentzian boosts corresponding to peculiar motion of the emitter and the observer (i.e. the Doppler effect).

Let us assume that light propagates in the outward direction. Using the standard formula for the Doppler redshift, we get in the static frame for the emitter

$$(\omega_{st})_{em} = (\omega_{GL})_{em} \sqrt{\frac{1 - v_{em}}{1 + v_{em}}}, \quad (41)$$

where subscripts “st” and “GL” indicate the static and GL frames, respectively. The same formula holds for the observer.

Then,

$$(1 + z)_{st} = \frac{(\omega_{st})_{em}}{(\omega_{st})_{obs}} = (1 + z)_{GL} \sqrt{\frac{1 - v_{em}}{1 + v_{em}}} \sqrt{\frac{1 + v_{obs}}{1 - v_{obs}}}. \quad (42)$$

Taking into account (30), we obtain

$$(1 + z)_{st} = \frac{\sqrt{1 - v_{obs}^2}}{\sqrt{1 - v_{em}^2}} = \frac{\sqrt{f_{obs}}}{\sqrt{f_{em}}}. \quad (43)$$

Here we reproduced the well-known result (see, e.g. Sec. 2.3.1 in [15]) which is ascribed usually to the gravitational time delay. Note, that this concept does not show up explicitly in the presented derivation, and we have only a combination of a “cosmological” redshift with the usual SR Doppler formula. If $r_{obs} > r_{em}$, light propagates outward, $f_{obs} > f_{em}$,

$z > 0$ and we have redshift. If $r_{obs} < r_{em}$, light propagates inward, $z < 0$ and the resulting effect is blueshift.

In the static coordinate system the interval of time t between two pulses remains the same during light propagation. As a result, the redshift is the ratio of values of the function relating proper time with the coordinate time t at the moments of emission and detection. In this situation it is possible to think that the time goes slowly in regions of large f . Moreover, if we imagine a space traveller who visits a vicinity of the event horizon and then returns, he/she will be younger than his twin staying on Earth. This is true even if the traveller has enough rocket fuel to move towards the black hole and back very slowly, so that it is hard to attribute the different time passed to usual Lorentz time dilation. However, it is necessary to keep in mind that such a kind of interpretation is connected with a particular coordinate system rather than with spacetime itself. In the Lemaitre coordinates, there is no gravitational time delay, and black hole twin paradox can be explained solely by SR Lorentz time dilation – to return back, the traveller must apply a huge Lorentz boost with respect to his “natural” free-falling frame.

The above formula (43) refers to a standard situation when both — an observer and an emitter, — are at rest in a static gravitational field. It is instructive to remind that there is an analogue of this situation in cosmology. This corresponds to the case of so-called tethered galaxies [16]. In this case the cosmological expansion is exactly compensated due to a peculiar velocity towards an observer, so the proper distance remains constant. However, the detected emission can be redshifted or blueshifted depending on a cosmological equation of state. This fact has been remarked only recently in [16] and shows clearly that the cosmological redshift can not be considered as a SR Doppler shift in an expanding Universe. In the case of black holes for free-falling observer and emitter, as we want to use the Lemaitre metric or its generalization, we can mimic this situation in the limit $r_{obs} \rightarrow \infty$. Then we can approximately consider that the observer is at rest. To have $r = const$, it is necessary for the emitter to have a peculiar velocity which exactly compensates the free-fall velocity. The emission will be always detected with a redshift. Usually this situation is considered in static coordinates, and it is interpreted in terms of gravitational time delay. However, we see that we can use a different interpretation, saying that in Lemaitre metric the non-zero redshift of an object at a constant proper distance has the same nature as non-zero redshifts of tethered galaxies. We also notice that the condition $r_{em} = const$ corresponds to

a constant light travel distance. So, an object with constant light travel distance is always redshifted — the property which is still valid in the FRW cosmology [17].

VIII. CONCLUSION

In this methodological note we proposed another interpretation of a redshift in spherically symmetric static spacetimes, including black holes. We split the calculation of the effect to two parts. 1) The first one includes transition to the synchronous system that is formed by free-falling particles. Here, the situation is very similar to that in cosmological FRW spacetime. 2) At the second step, we return to the static system by means of the local Lorentz boost. This enables us to consider it as a kind of the Doppler shift and apply well-known corresponding formulas. It is worth noting that calculations on step 1 are realized by us by two methods. We found (i) the frequency from equations of motion, (ii) the wavelength taking into account that different points move with different velocities (in analogy with cosmology). Thus, we see that there is no crucial difference between interpretations of the redshift in cosmological and black hole spacetimes. Rather, the situation in the latter case can be thought of as a combination of the former one and the Doppler effect.

The obtained results have a rather curious consequence. As it is noticed above, an observer crossing the horizon registers a signal from infinity with $z = 1$. This allows the observer to identify the moment of crossing the event horizon. In doing so, an observer should use information about the photon emitted at infinity whereas he is unable to do this using only local measurements. (It is instructive to remind that formally another kind of pure local measurements can in principle locate the moment of the horizon crossing but that method is rather sophisticated and does not operate with entities having direct physics meaning [18].)

When speaking about the cosmological spacetimes throughout the paper, we implied the FRW spacetimes as opposed to the static ones like, say, the Schwarzschild metric. For completeness, we have to mention that there exist also special cases when cosmological spacetimes themselves admit presentation in a static form. This includes the de Sitter and Milne solutions when both forms (explicitly static and pure cosmological) are possible. In [19] six cosmological metrics with stationary frames have been listed, and corresponding redshifts studied. It was shown that in all these cases the cosmological redshift can be

represented as a combination of SR Doppler shift and gravitational time delay, and, so it cannot be considered as a “fundamental” shift. In our paper we did just the opposite, showing that gravitational redshift in a stationary frame can be represented as a combination of the Doppler shift and a “cosmological” redshift. This means that the question of what type of redshift should be considered as a more fundamental one is the matter of convenience. On the other hand, we can argue that since a synchronous system can be constructed for any spacetime and static systems exist only for some particular metrics, the “cosmological” description of a redshift presented in our paper has more general significance.

Acknowledgments

This work was funded by the subsidy allocated to Kazan Federal University for the state assignment in the sphere of scientific activities. SBP acknowledges support from the Russian Science Foundation (grant 14-12-00146).

-
- [1] E. Harrison, *Cosmology: The Science of the Universe*. The 2nd edition. Cambridge University Press, 2000.
 - [2] J. V. Narlikar. Spectral shifts in general relativity. *Am. J. Phys.* 62, 903, (1994).
 - [3] J. Synge. *Relativity: the general theory*, North-Holland, Amsterdam (1960).
 - [4] A. Radosz, A.T. Augousti, K. Ostasiewicz, The Doppler shift in a Schwarzschild spacetime. *Physics Letters A* 373, 801 (2009).
 - [5] C. W. Misner, K. S. Thorne, J. A. Wheeler. *Gravitation*. W. H. Freeman and company. San Francisco (1973).
 - [6] A. Liddle. *An Introduction into modern cosmology*. Third Edition. John Wiley & Sons, Ltd., 2015.
 - [7] A. Toporensky and O. Zaslavskii, Redshift of a photon emitted along the black hole horizon, *Europ. Phys. J. C* [arXiv:1611.09807].
 - [8] L. D. Landau and E. M. Lifshitz, *The Classical Theory of Fields* Pergamon Press, New York (1971).
 - [9] K. Martel and E. Poisson, Regular coordinate systems for Schwarzschild and other spherical

- spacetimes, *Am. J. Phys.* 69, 476 (2001); [arXiv:gr-qc/0001069].
- [10] A. J. S. Hamilton, J. P. Lisle, The river model of black holes, *Am.J.Phys.* 76, 519 (2008) [arXiv:gr-qc/0411060].
 - [11] A. Radosz, A. T. Augousti, A. Siwek, On the nature of cosmological redshift and spectral shift in Schwarzschild-like and other spacetimes, *Gen. Rel. Grav.* v. 45, pp. 705-715 (2013).
 - [12] K. Kassner, Why ghosts don't touch: a tale of two adventurers falling one after another into a black hole, *Eur. J. Phys.* 38, 015605 (2017) [arXiv:1608.07511].
 - [13] A. Toporensky and S. Popov, The Hubble flow: an observer's perspective, *Phys.Usp.* 57, 708-713 (2014) (*Usp.Fiz.Nauk* 184, 767 (2014)) [arXiv:1311.2472].
 - [14] G. Horndeski, Conservation of Charge and the Einstein-Maxwell Field Equations, *J.Math.Phys.* 17 (1976) 1980.
 - [15] V. P. Frolov and I. D. Novikov, *Physics of black holes*, Kluwer Academic Publishers, Dordrecht, Boston, London (1998).
 - [16] Tamara M. Davis, Charles H. Lineweaver, John K. Webb, Solutions to the tethered galaxy problem in an expanding universe and the observation of receding blueshifted objects, *Am.J.Phys.* 71:358-364, 2003 [arXiv:astro-ph/0104349].
 - [17] A.V. Toporensky, S.B. Popov, Cosmological redshift, recession velocities and acceleration measures in FRW cosmologies, *Astron. and Astrophys. Trans.* v. 29, pp. 65 - 88 (2015) . [arXiv:1503.05147].
 - [18] I. V. Tanatarov and O. B. Zaslavskii, What happens to Petrov classification, on horizons of axisymmetric dirty black holes, *Journ. of Math. Phys.* 55, 022502 (2014) [arXiv:1211.4376].
 - [19] F. Melia, Cosmological Redshift in FRW Metrics with Constant Spacetime Curvature, *MNRAS* 422, 1418 (2012) [arXiv:1202.0775].